

Distribution and molecular analysis of *Blastocystis* subtypes from gastrointestinal symptomatic and asymptomatic patients in Iran

Asma Delshad¹, Mehrzad Saraei^{2,3}, Safar Ali Alizadeh^{4,5}, Soheila Rahimi Niaraki¹,
Mahmood Alipour⁶, Bahram Hosseinbigi⁴, Arezoo Bozorgomid⁷, Elham Hajjalilo^{2,3}

1. Student Research Committee, Qazvin University of Medical Sciences, Qazvin, Iran.
2. Cellular and Molecular Research Center, Research Institute for Prevention of Non- Communicable Disease, Qazvin University of Medical Sciences, Qazvin, Iran.
3. Department of Parasitology and Mycology, Qazvin University of Medical Sciences, Qazvin, Iran.
4. Medical Microbiology Research Center, Qazvin University of Medical Sciences, Qazvin, Iran.
5. Department of Microbiology, Qazvin University of Medical Sciences, Qazvin, Iran.
6. Department of Social Medicine, Qazvin University of Medical Sciences, Qazvin, Iran.
7. Infectious Diseases Research Center, Health Institute, Kermanshah University of Medical Sciences, Kermanshah, Iran

Abstract

Introduction: *Blastocystis* is a common intestinal parasite of human and animal hosts. The parasite has 17 subtypes, and among those at least nine subtypes (ST1-ST9) are found in human hosts.

Objective: The aim of the present study was to investigate the presence of different subtypes of *Blastocystis* spp. among the patients referred to Velayat hospital of Qazvin province, Iran.

Methods: Overall, 864 stool samples were examined by using formalin-ethyl acetate concentration method and Trichrome staining. All specimens were cultured in clotted fetal bovine medium. Later, DNA extraction and PCR amplification of 18S ribosomal RNA gene region was conducted and phylogenetic tree constructed.

Results: The results revealed 7.9% (68/864) of the study population were infected with *Blastocystis*. Intestinal symptoms were observed in 61% (36/59) of individuals positive for *Blastocystis*, with abdominal pain in 58% (21/36) of cases which was more frequent than other intestinal signs. No significant relationship was observed among the study variables. By molecular and phylogenetic analysis, three subtypes ST1 (45%), ST2 (30%) and ST3 (23%) of parasite were identified.

Conclusion: This study showed ST1 subtype was the predominant subtype among the positive specimens, meanwhile the highest haplotype and nucleotide diversity were clarified in ST3 subtype.

Keywords: *Blastocystis*, subtype, Phylogenetic analysis, Iran.

DOI: <https://dx.doi.org/10.4314/ahs.v20i3.21>

Cite as: Delshad A, Saraei M, Alizadeh SA, Niaraki SR, Alipour M, Hosseinbigi B, et al. Distribution and molecular analysis of *Blastocystis* subtypes from gastrointestinal symptomatic and asymptomatic patients in Iran. *Afri Health Sci.*2020;20(3): 1179-1189. <https://dx.doi.org/10.4314/ahs.v20i3.21>

Corresponding author:

Mehrzad Saraei,
Department of Parasitology and Mycology,
Qazvin University of Medical Sciences, Qazvin, Iran
E-mail address: msaraei@qums.ac.ir
and
Elham Hajjalilo,
Department of Parasitology and Mycology,
Qazvin University of Medical Sciences, Qazvin, Iran
E-mail address: e.hajjalilo@qums.ac.ir

Introduction

Blastocystis is an anaerobic parasite of human and animal intestinal tract. According to phylogenetic approach, the parasite belongs to Stramenopiles of the Eukaryota¹. The parasite is a common intestinal microorganism reported in epidemiological surveys². High frequency of this parasite may be due to the reluctance of physicians to treat self-limiting infections or resistance of the parasite to anti-parasitic drugs in conventional treatment, therefore, *Blastocystis* could easily colonize intestinal empty niches. Moreover, *Blastocystis* could reside as a non-pathogenic microorganism in the healthy human